

CETUP* 2024

Monday, June 17, 2024 - Friday, July 19, 2024

Lead/Deadwood Middle School



THE INSTITUTE
FOR UNDERGROUND SCIENCE
AT SURF

Book of Abstracts

Contents

Charged-current (anti)neutrino-nucleon scattering and QED nuclear medium effects . . .	1
New Directions for ALP Searches Combining Nuclear Reactors and Haloscopes	1
CP-Violation with Neutrino Disappearance	1
Dark matter annihilation signals from Sagittarius analogues	2
Dark matter at the high mass frontier	2
New Constraints on Neutrino-Dark Matter Interactions	2
Gravitational Wave Symphony from Oscillating Spectator Scalar Fields	3
Neutrinos from the Sun can discover dark matter-electron scattering	3
TALK: Dark Radiation Isocurvature from Cosmological Phase Transitions	4
Welcome by Mike Headley, Executive Director, South Dakota Science and Technology Authority/SURF	4
The Institute for Underground Science at SURF Overview and Update (room 132)	4
Discussion, Collaborations and Lunch	4
TALK: Dark matter at the high mass frontier	4
Discussion and Collaboration	5
Shuttle Pickup for Underground Tour - Travel to SURF Yates Admin, 2nd Floor Vault . . .	5
SURF Overview by Mike Headley, Executive Director, South Dakota Science and Technology Authority/Laboratory Director, SURF	5
SURF Underground Tour Safety Training, Waivers, Lunch at 11:00am, PPE, Shuttle to Ross	5
Shuttle, PPE, Return to Middle School	5
TALK: Dark Matter Annihilation Signals from Sagittarius Analogues	5
TALK: (Speaker: Pearl Sandick)	6
Discussions, Collaboration, and Lunch	6

TALK: TBD (open slot available)	6
Discussions and Collaboration	6
Constraining Dwarf Galaxy Dark Matter Distributions: Spherical Jeans Analyses for Line-of-Sight and 3D Velocity Data	6
TALK: TBD (open slot available)	6
Discussions and Collaboration	7
Travel to Ethnobotanical Garden at SURF (via shuttle)	7
Guest Speaker at the Ethnobotanical Gardens (Speaker: Rochelle Zens)	7
Travel to Middle School	7
Discussions and Collaboration	7
TALK: Analytic Results on Dark Matter Velocity Distributions	7
TALK: Probing Neutrinophilic Dark Matter: From Colliders to Supernovae	7
Discussion, Collaboration, and Lunch	8
TALK: Decoding the Mystery of Dark Matter with Celestial Objects (Speaker: Anupam Ray)	8
TALK: Thermal Axion and Dark Radiation	8
TALK: The Power of the Dark Sink	8
Discussions, Collaboration, and Lunch	9
TALK: Jet-SIFTing for Dark Sector Physics in a Hidden Valley	9
Discussion and Collaboration	9
TALK: SPLENDOR: Narrow-gap Quantum Materials for Light Dark Matter	9
TALK: SPLENDOR: narrow-gap quantum materials for light dark matter	10
Shuttle to Middle School for Discussion and Collaboration	10
TALK: TBD	10
Discussion and Collaboration	10
TALK: Dynamical Generation of the Baryon Asymmetry from a Scale Hierarchy	10
TALK: Gauged global strings	10
Discussion, Collaboration, and Lunch	11
TALK: Cosmic Stability of Dark Matter from Pauli Blocking	11
Discussion and Collaboration	11

TALK: BSM Physics with Gravitational Wave Detectors	11
Discussion, Collaboration, and Lunch	12
TALK: Cosmological Stasis and Its Realization from Dynamical Scalars	12
TALK: Updates on the Migdal effect and hydrogen doping for dark matter detection	12
Discussion and Collaboration	13
TALK: Dark Matter Raining on DUNE and Other Large Volume Detectors	13
TALK: Gravitational Waves from Cosmic Superstrings and Gauge Strings	13
Discussion, Collaboration, and Lunch	14
TALK: Introduction to DAMSA, A Novel Dark Matter Search Experiment at an Accelerator	14
Discussion and Collaboration	14
TALK: Dark Matter in the Time of Gravitational Waves	14
TALK: Probing self-interacting sterile neutrino dark matter with the diffuse supernova neu- trino background	14
Discussion, Collaboration, and Lunch	15
TALK: Introduction to DAMSA, A Novel Dark Matter Search Experiment at an Accelerator	15
Discussion and Collaboration	15
TALK: Journey with Sourced Millicharged Particles and Dark Photons	15
TALK: Asymptotically Free E_6 GUT and the Generation of Neutrino Mass	16
Discussion, Collaboration, and Lunch	16
TALK: TALK: Gravitational Wave Symphony from Oscillating Spectator Scalar Fields	16
Discussion and Collaboration	16
TALK: Probing Baryon Number Violation in Neutron Stars	16
TALK: TBD (open slot available)	17
Discussion, Collaboration, and Lunch	17
TALK: TBD (open slot available)	17
Discussion and Collaboration	17
TALK: Does the Sun Have a Dark Disk?	17
TALK: TBD (open slot available)	18
Discussion, Collaboration, and Lunch	18

Independence Day Festivities in Lead	18
TBD (open slot available)	18
TALK: Neutrino Interactions	18
Discussion, Collaboration, and Lunch	18
TALK: Title TBD (Speaker: Volodymyr Takhistov)	18
Discussion and Collaboration	19
The Institute for Underground Science at SURF Update	19
TALK: CP - Violation with Neutrino Disappearance	19
Discussion, Collaboration, and Lunch	19
TALK: A 17 MeV pseudoscalar and the MiniBooNE, LSND and ATOMKI anomalies	19
Discussion and Collaboration	20
TALK: Charged-current (anti)neutrino-nucleon Scattering and QED Nuclear Medium	20
TALK: Constraining Non-standard Neutrino Interactions with Neutral Current Events of Long-Baseline Oscillation Experiments	20
Discussion, Collaboration, and Lunch	21
TALK: A SURF Low Background Module (SLoMo)	21
Discussion and Collaboration	21
Shuttle to Middle School	21
Shuttle to Sanford Lab Visitor Center	21
TALK: Probing Heavy Neutral Lepton on Muon Collider and Beamdump Experiments	21
Discussion, Collaboration, and Lunch	22
Shuttle to Ethnobotanical Garden	22
Guest Speaker at Ethnobotanical Garden - Rochelle Zens	22
Shuttle to Middle School	22
TALK: Neutrino and light dark matter physics with directional detectors	22
Discussion and Collaboration	23
TALK: Title TBD (Speaker: Garv Chauhan)	23
TALK: The Scotos Side of Neutrinos	23
Discussion, Collaboration, and Lunch	23
TALK: Does the Sun have a dark disk?	23

Discussion and Collaboration	24
Coffee/Social with Students	24
TALK: Direct Collapse Black Holes from Dark Matter Annihilation	24
TALK: Quasi (pseudo)-Dirac Neutrinos	24
Discussion and Collaboration	24
TALK: Neutrinos From the Sun Can Discover Dark Matter-electron Scattering	24
TALK: E6 Unification: Intermediate Symmetries, Fermion Masses and Proton Decay	25
Discussion, Collaboration, and Lunch	25
TALK: TBD (open slot available)	25
Discussion and Collaboration	25
TALK: New Directions for ALP Searches Combining Nuclear Reactors and Haloscopes	25
TALK: Decoding the Flavor Evolution of Supernova Neutrinos: Shedding Light on Neutrino Electromagnetic Properties and Supernova Dynamics	26
Discussion, Collaboration, and Lunch	26
TALK: TBD (open slot available)	26
Discussion and Collaboration	26
TALK: Probing SUSY at Gravitational Wave Observations	26
TALK: Embedding the Universal See-Saw Mechanism in Pati-Salam Model	27
Discussion, Collaboration, and Lunch	27
TALK: eV Sterile Neutrino Paradox	27
Discussion and Collaboration	28
Shuttle Pickup at Hampton Inn, travel to Sanford Lab Homestake Visitor Center	28
Shuttle Pickup at Middle School, Travel to E&O Building	28
SURF Overview by Mike Headley, Executive Director, South Dakota Science and Technol- ogy Authority	28
SURF Underground Tour Safety, Waivers, Lunch at 11:00 w/ Cultural Awareness Commit- tee, PPE and Shuttle to Ross	28
Shuttle, PPE, Return to Hampton Inn	28
TALK: TBD (open slot available)	28
TALK: Skimming tau neutrinos and optical Cherenkov signals	29
Discussion, Collaboration, and Lunch	29

TALK: TBD (open slot available)	29
Discussion and Collaboration	29
Road Scholar Group - CETUP* Guest Speaker (Mary Hall Reno)	29
Dynamical Generation of the Baryon Asymmetry from a Scale Hierarchy	29
Discussion and Collaboration	30
The Power of the Dark Sink	30
Analytic Results on Dark Matter Velocity Distributions	30
Probing SUSY at Gravitational Wave Observatories	31
-TBD-	31
Gravitational waves from cosmic superstrings and gauge strings	31
Journey with Sourced Millicharged Particles and Dark Photons	31
gauged global strings	32
Probing Baryon Number Violation in Neutron Stars	32
Welcome by Mike Headley, Executive Director, South Dakota Science and Technology Authority/SURF	32
Shuttle to Hampton Inn	32
Modular Flavor Symmetry and Neutrino Oscillations	33
SPLENDOR: narrow-gap quantum materials for light dark matter	33
Updates on the Migdal effect and hydrogen doping for dark matter detection	33
Dark Radiation Isocurvature from Cosmological Phase Transitions	34
The Scotos side of Neutrinos	34
Does the Sun Have a Dark Disk	34
Probing Heavy Neutral Lepton at Muon Collider and Beamdump Experiments	35
Constraining non-standard neutrino interactions with neutral current events at long-baseline oscillation experiments	35
Embedding the Universal See-Saw Mechanism in Pati-Salam Model	36
Asymptotically free E_6 GUT and the generation of neutrino mass	36
Thermal Axion and Dark Radiation	36
E6 unification: intermediate symmetries, fermion masses and proton decay	37
Constraining Dwarf Galaxy Dark Matter Distributions: Spherical Jeans Analyses for Line-of-Sight and 3D Velocity Data	37

-	37
Probing Neutrinophilic Dark Matter: From Colliders to Supernovae	38
Neutrino Interactions/TBD	38
Decoding the Flavor Evolution of Supernova Neutrinos: Shedding Light on Neutrino Elec- tromagnetic Properties and Supernova Dynamics	38
TBD	39
Quasi (Pseudo)-Dirac neutrinos	39
Dark Matter Raining on DUNE and Other Large Volume Detectors	39
Cosmological Stasis and Its Realization from Dynamical Scalars	40
Neutrino and light dark matter physics with directional detectors	40
Direct Collapse Black Holes from Dark Matter Annihilation	41
Probing self-interacting sterile neutrino dark matter with the diffuse supernova neutrino background	41
Planting Native Plants at Sanford Lab Homestake Visitor Center	42
Communicating with Media (Speakers: Ann Melti and Mike Ray, SDSTA/SURF)	42
Communicating with Media - 3 Minute Challenge (Ann Melti and Mike Ray - SDSTA/SURF)	42
Career Panel	42
Scientist Mentor/Student Speed Match Up and Shadow	42
Communicating with Media - Ann Melti and Mike Ray (SDSTA/SURF)	42
3 Minute Challenge - Ann Melti and Mike Ray (SDSTA/SURF)	42
Jet-SIFTing for Dark Sector Physics in a Hidden Valley	43
Skimming tau neutrinos and optical Cherenkov signals	43
Detection of cosmogenic (B)SM signals with nuclear inelastic scattering	43
BSM Physics with Gravitational Wave Detectors	43
eV Sterile Neutrino Paradox	44
CETUP* Workshop Overview	44
Discussion and Collaboration	44
TALK: New Constraints on Neutrino-Dark Matter Interactions	44
TALK: TBD (open slot available)	45
Shuttle, Travel to Ross	45

Does the Sun have a dark disk?	45
TALK: Direct Collapse Black Holes from Dark Matter Annihilation	45
TALK: TBD (open slot available)	46
Discussions, Collaboration and Lunch	46
Current Challenges and Opportunities in Physics Community (open discussion)	46
Global Scientist Social	46
SURF Underground Tour	46
Discussion and Collaboration; Mentor/Student Shadow	46
A SURF Low Background Module (SLoMo)	47
The Institute for Underground Science at SURF Overview and Update (room 132)	47
The Institute for Underground Science at SURF Overview and Update (room 132)	47
The Institute for Underground Science at SURF Overview and Update (room 132)	47
REMINDER NOTE ONLY: please make sure to bring a brown bag lunch for today	47
TALK: Detection of cosmogenic (B)SM signals with nuclear inelastic scattering	47

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 77

Charged-current (anti)neutrino-nucleon scattering and QED nuclear medium effects

Author: Oleksandr Tomalak¹

¹ *Los Alamos National Laboratory*

Corresponding Author: tomalak@lanl.gov

Charged-current quasielastic neutrino scattering is the signal process in neutrino oscillation experiments and requires precise theoretical prediction for the analysis of modern and future experimental data, starting with the nucleon axial-vector form factor. In this talk, I compare a new MINERvA measurement of this form factor with lattice-QCD calculations and deuterium bubble-chamber data, provide uncertainty projections for future extractions, present recent calculations of radiative corrections to charged-current processes, and investigate the potential of neutrino scattering data on constraining nucleon- and quark-level interactions beyond the Standard Model.

The exchange of photons with nuclear medium modifies (anti)neutrino and electron scattering cross sections. We study the distortion of (anti)neutrino-nucleus and charged lepton-nucleus cross sections, medium-induced bremsstrahlung, and estimate the QED-medium effects on the final-state kinematics and scattering cross sections. We find new permille-to-percent level effects, which were never accounted for in either (anti)neutrino-nucleus or electron-nucleus scattering. We quantitatively compute the effects of Glauber photon-mediated multiple re-scattering within the nuclear medium and find that the relativistic charged lepton acquires a momentum of order 10 MeV transverse to its direction of propagation inside the nucleus. This broadening sizably deflects expected electron and muon tracks and suppresses scattering cross sections. Precise extraction of the nucleon and nuclear structure by electron and muon probes should, thus, take the QED nuclear medium angular redistribution of particles into account.

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 78

New Directions for ALP Searches Combining Nuclear Reactors and Haloscopes

Author: Vedran Brdar¹

¹ *Oklahoma State University*

Corresponding Author: vedran.brdar@okstate.edu

I will discuss a recently proposed novel experimental setup for axion-like particle (ALP) searches. Nuclear reactors produce a copious number of photons, a fraction of which could convert into ALPs via the Primakoff process in the reactor core. The generated flux of ALPs leaves the nuclear power plant, and its passage through a region with a strong magnetic field results in efficient conversion to photons, which can be detected. Such a magnetic field is the key component of axion haloscope experiments. I will discuss existing setups featuring an adjacent nuclear reactor and axion haloscope and I will demonstrate that the obtained sensitivity projections complement constraints from existing laboratory experiments, e.g., light-shining-through-walls.

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 79

CP-Violation with Neutrino Disappearance

Author: Peter Denton¹

¹ *Brookhaven National Laboratory*

Corresponding Author: peterbd1@gmail.com

The best way to probe CP violation in the lepton sector is with long-baseline accelerator neutrino experiments in the appearance mode: the appearance of ν_e in predominantly ν_μ beams. Here we show that it is possible to discover CP violation with disappearance experiments only, by combining JUNO for electron neutrinos and DUNE or Hyper-Kamiokande for muon neutrinos. While the maximum sensitivity to discover CP is quite modest (1.6σ with 6 years of JUNO and 13 years of DUNE), some values of δ may be disfavored by $> 3\sigma$ depending on the true value of δ .

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 80

Dark matter annihilation signals from Sagittarius analogues

Author: Nassim Bozorgnia¹

¹ *University of Alberta*

Corresponding Author: nbozorgnia@ualberta.ca

Dwarf spheroidal galaxies such as Sagittarius are dark matter dominated, and therefore unique candidates for indirect dark matter searches. In order to accurately predict the dark matter annihilation signal from dwarf spheroidal galaxies, it is crucial to correctly model the phase space distribution of dark matter in them. Hydrodynamical simulations of galaxy formation provide important information on the dark matter distribution in dwarf spheroidal analogues. I will present the dark matter density profile and velocity distribution of the Sagittarius dwarf spheroidal galaxy extracted from state-of-the-art hydrodynamical simulations. In addition to the annihilation signals from dark matter particles bound to Sagittarius, we consider for the first time the annihilation of dark matter particles bound to the Milky Way that overlap spatially with Sagittarius. I will discuss the implications of this dark matter population for velocity-dependent dark matter annihilation models.

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 81

Dark matter at the high mass frontier

Author: Joe Bramante¹

¹ *Queen's University*

Corresponding Author: joseph.bramante@queensu.ca

A number of theories predict that dark matter is a supermassive particle or composite state. Discovering dark matter in this high mass regime requires different approaches. I will survey recent developments, including composite dark matter that produces unique signatures in underground experiments and dark matter detectable through thermonuclear reactions in Antarctic ice. I will also survey certain experiments from the 80s and 90s, which still provide the best sensitivity to many varieties of high mass dark matter.

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 82

New Constraints on Neutrino-Dark Matter Interactions

Author: Bhupal Dev¹

¹ *Washington University in St. Louis*

Corresponding Author: bdev@wustl.edu

We present a comprehensive analysis of nonstandard neutrino interactions with the dark sector in an effective field theory framework. We implement a full catalog of constraints on the parameter space of the neutrino-dark matter/mediator couplings and masses, including bounds coming from cosmology, astrophysics, as well as new laboratory constraints, such as from invisible Z decays and rare meson decays. We find that most of the benchmarks in the dark matter mass-coupling plane adopted in previous studies to get an observable effect are actually ruled out by a combination of these constraints. Finally, as an application of our results, we consider the case of galactic supernova neutrinos, identify new benchmark points for future observational prospects of the attenuation of the galactic supernova neutrino flux, compute the full set of cascade equations and sky maps for different dark matter density profiles in the Galaxy, and comment on their implications for the detection prospects in future large-volume neutrino experiments such as DUNE, Hyper-K and JUNO.

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 83

Gravitational Wave Symphony from Oscillating Spectator Scalar Fields

Author: Yanou Cui^{None}

Corresponding Author: yanou.cui@ucr.edu

Spectator scalar fields can be generically present in the early Universe, and are potentially dark matter candidates. We investigate the prospect of such scalars loop as a generic source of stochastic gravitational wave background (SGWB) due to parametric resonance during their oscillation phase. By systematically analyzing benchmark models through lattice simulations and considering a wide range of parameters, we demonstrate that such a scenario can lead to detectable signals in GW detectors over a broad frequency range and potentially address the recent findings by Pulsar Timing Array experiments. Furthermore, we show that these models naturally yield viable ultra-light, wave-like dark matter candidates and/or dark radiation detectable by CMB observatories. We also explore the potential for realizing low-scale baryogenesis from such an oscillating scalar system and the resultant detectable SGWB signals. This study highlights an intriguing connection between the stochastic gravitational wave background (SGWB) and terrestrial probes for low-energy new particle physics.

Request to give talk on Wednesday, July 3 in the afternoon

CETUP 2024 - schedule of discussions, collaborations, and tours to be determined / 84

Neutrinos from the Sun can discover dark matter-electron scattering

Author: Ranjan Laha¹

¹ *Indian Institute of Science, Bengaluru, India*

Corresponding Author: ranjanlaha@iisc.ac.in

We probe dark matter-electron scattering using high-energy neutrino observations from the Sun. Dark matter (DM) interacting with electrons can get captured inside the Sun. These captured DM may annihilate to produce different Standard Model (SM) particles. Neutrinos produced from these SM states can be observed in IceCube and DeepCore. Although there is no excess of neutrinos from the Solar direction, we find that the current data-sets of IceCube and DeepCore set the strongest constraint on DM-electron scattering cross section in the DM mass range 10 GeV to 10^5 GeV. Our

work implies that future observations of the Sun by neutrino telescopes have the potential to discover DM-electron interaction.

85

TALK: Dark Radiation Isocurvature from Cosmological Phase Transitions

Corresponding Author: peizhi.du@rutgers.edu

Speaker: Peizhi Du (Rutgers University)

Authors: Matthew Buckley, Peizhi Du, Nicolas Fernandez, Mitchell Weikert

Cosmological first-order phase transitions are typically associated with physics beyond the Standard Model, and thus of great theoretical and observational interest. In this talk, I will show that a broad class of non-thermal first-order phase transitions could generate distinct large-scale isocurvature in dark radiation that can be observable in the CMB. We derive constraints on ___ from phase transitions based on CMB+BAO data, which can be much stronger than that from adiabatic initial conditions. I will also demonstrate that since perturbations of dark radiation have a non-Gaussian origin, searches for non-Gaussianity in the CMB could also place a stringent bound on ___.

86

Welcome by Mike Headley, Executive Director, South Dakota Science and Technology Authority/SURF

87

The Institute for Underground Science at SURF Overview and Update (room 132)

88

Discussion, Collaborations and Lunch

89

TALK: Dark matter at the high mass frontier

Corresponding Author: joseph.bramante@queensu.ca

Author: Joe Bramante

A number of theories predict that dark matter is a supermassive particle or composite state. Discovering dark matter in this high mass regime requires different approaches. I will survey recent developments, including composite dark matter that produces unique signatures in underground experiments and dark matter detectable through thermonuclear reactions in Antarctic ice. I will also survey certain experiments from the 80s and 90s, which still provide the best sensitivity to many varieties of high mass dark matter.

90

Discussion and Collaboration

91

Shuttle Pickup for Underground Tour - Travel to SURF Yates Admin, 2nd Floor Vault

Pick up at the Middle School

This shuttle is only for the CETUP* participants that are going on the underground tour. Other participants not going on the tour will be able to stay at the Middle School and continue to work and carry on discussions.

92

SURF Overview by Mike Headley, Executive Director, South Dakota Science and Technology Authority/Laboratory Director, SURF

93

SURF Underground Tour Safety Training, Waivers, Lunch at 11:00am, PPE, Shuttle to Ross

Corresponding Authors: lbaatz@sanfordlab.org, jmolina@sanfordlab.org

Only the CETUP* participants that are going on the underground tour will attend. Other participants not going on the tour will be able to stay at the Middle School and continue to work and carry on discussions.

94

Shuttle, PPE, Return to Middle School

95

TALK: Dark Matter Annihilation Signals from Sagittarius Analogues

Corresponding Author: nbozorgnia@ualberta.ca

Author: Nassim Bozorgnia

Dwarf spheroidal galaxies such as Sagittarius are dark matter dominated, and therefore unique candidates for indirect dark matter searches. In order to accurately predict the dark matter annihilation signal from dwarf spheroidal galaxies, it is crucial to correctly model the phase space distribution of

dark matter in them. Hydrodynamical simulations of galaxy formation provide important information on the dark matter distribution in dwarf spheroidal analogues. I will present the dark matter density profile and velocity distribution of the Sagittarius dwarf spheroidal galaxy extracted from state-of-the-art hydrodynamical simulations. In addition to the annihilation signals from dark matter particles bound to Sagittarius, we consider for the first time the annihilation of dark matter particles bound to the Milky Way that overlap spatially with Sagittarius. I will discuss the implications of this dark matter population for velocity-dependent dark matter annihilation models.

96

TALK: (Speaker: Pearl Sandick)

97

Discussions, Collaboration, and Lunch

98

TALK: TBD (open slot available)

99

Discussions and Collaboration

100

Constraining Dwarf Galaxy Dark Matter Distributions: Spherical Jeans Analyses for Line-of-Sight and 3D Velocity Data**Corresponding Author:** isgoldstein@tamu.edu

Author: Isabelle Goldstein

Co-author: Louie Strigari

The stellar kinematics in dwarf galaxies can provide a wealth of information about its underlying dark matter distribution. Using line of sight velocity measurements for six classical dwarf galaxies of the Milky Way, we study whether ultralight bosonic dark matter is consistent with the gravitational potential extracted from stellar kinematics. It shows that axion-like particles with masses of order _____ eV are inconsistent with the potential distribution in classical dwarf galaxies unless the hierarchical assembly of the Milky Way did not trace the mean evolution of Milky Way size halos. We also explore the use of three dimensional velocity measurements from Gaia data in spherical Jeans analyses to further constrain dark matter distributions.

101

TALK: TBD (open slot available)

102

Discussions and Collaboration

103

Travel to Ethnobotanical Garden at SURF (via shuttle)

104

Guest Speaker at the Ethnobotanical Gardens (Speaker: Rochelle Zens)

Corresponding Author: rzens@sanfordlab.org

105

Travel to Middle School

106

Discussions and Collaboration

107

TALK: Analytic Results on Dark Matter Velocity Distributions

Corresponding Author: jkumar@hawaii.edu

Author: Jason Kumar

We review recent work on determining dark matter velocity distributions from analytic methods of classical mechanics, and the potential impact of these results on dark matter indirect detection strategies. In particular, we discuss velocity-dependent dark matter annihilation in subhalos and in the Galactic Center. We compare the results of these analytic methods to those obtained in large N numerical simulations, and discuss future work.

108

TALK: Probing Neutrinophilic Dark Matter: From Colliders to Supernovae

Corresponding Author: dtuckler@triumf.ca

Author: Douglas Tuckler

New beyond-the-Standard Model mediators that couple predominantly to neutrinos are not yet probed by existing experimental searches. Such a neutrinophilic mediator is well motivated for addressing the origin of several neutrino-portal dark matter candidates, including thermal freeze-out and sterile-neutrino dark matter scenarios. In this talk, we explore the sensitivity to this scenario from two different approaches. In the first part of the talk we will explore the potential of the Forward Physics Facility (FPF) using the so-called “mono-neutrino signature”: neutrino charged-current scattering events associated with large missing transverse momentum, and excessive apparent tau-neutrino events. We will show that with this smoking-gun signature, the FPF has excellent sensitivity to probe this model in new regions of parameter space. In the second part of the talk we focus on astrophysical constraints on sterile neutrino DM, namely from core-collapse supernovae. Production and emission of DM can result in excessive energy loss of the supernova, leading to additional cooling. We will show that supernova cooling can constrain new regions of parameter space that complements terrestrial and cosmological probes. The results of this work present a nice complementarity among the collider and astrophysical frontiers.

109

Discussion, Collaboration, and Lunch

110

TALK: Decoding the Mystery of Dark Matter with Celestial Objects (Speaker: Anupam Ray)

Corresponding Author: anupam.ray@berkeley.edu

Dark Matter (DM) remains mysterious. Despite decades of experimental and theoretical efforts, its microscopic identity is still unknown to us. In this talk, I will walk you through how a variety of celestial objects can be utilised as powerful DM detectors. This astrophysical probe, complementary to the terrestrial and cosmological probes, covers a significant portion of the DM parameters (DM mass and its interaction strength with nucleons) which are otherwise remains elusive.

111

TALK: Thermal Axion and Dark Radiation

Corresponding Author: fazlollah.hajkarim@ou.edu

Author: Fazlollah Hajkarim

We provide a comprehensive analysis of thermal axion production in the early universe, focusing on the KSVZ and DFSZ models. We extend our calculations to incorporate multiple mass thresholds, including the QCD phase transition, and provide a continuous production rate across these regimes. Employing updated cosmological data, we refine constraints on axion mass and examine the implications for dark radiation quantified as an effective number of additional neutrino species (ΔN_{eff}). Our rigorous approach revisits traditional approximations in axion production, highlighting the importance of precise calculations in anticipation of future CMB surveys and large scale structure observations. Also, we examine the validity of different computational approaches for the calculation of effective number of additional neutrino species.

112

TALK: The Power of the Dark Sink

Corresponding Author: rmcgehee@umn.edu

Author: Robert McGehee

We describe a simple dark sector structure which, if present, has implications for the direct detection of dark matter (DM): the Dark Sink. A Dark Sink transports energy density from the DM into light dark-sector states that do not appreciably contribute to the DM density. As an example, we consider a light, neutral fermion which interacts solely with DM via the exchange of a heavy scalar. We illustrate the impact of a Dark Sink by adding one to a DM freeze-in model in which couples to a light dark photon

which kinetically mixes with the Standard Model (SM) photon. This freeze-in model (absent the sink) is itself a benchmark for ongoing experiments. In some cases, the literature for this benchmark has contained errors; we correct the predictions and provide them as a public code. We then analyze how the Dark Sink modifies this benchmark, solving coupled Boltzmann equations for the dark-sector energy density and DM yield. We check the contribution of the Dark Sink's to dark radiation; consistency with existing data limits the maximum attainable cross section. For DM with a mass between , adding the Dark Sink can increase predictions for the direct detection cross section all the way up to the current limits.

113

Discussions, Collaboration, and Lunch

114

TALK: Jet-SIFTing for Dark Sector Physics in a Hidden Valley

Corresponding Author: jww004@shsu.edu

Author: Joel Walker

We consider a Hidden Valley model which generates showering from strong dynamics within the dark sector followed by decays back into Standard Model states. Our interest is the limit of smaller dark pion masses, which create a high multiplicity of final states. The reconstruction of dark sector masses in such a setting is obscured by a thick combinatoric background. We apply the new SIFT (Scale-Invariant Filtered Tree) jet clustering algorithm to the reconstruction of simulated events of this type. By cutting an ordered slice through possible recombinations, the SIFT algorithm may help lift backgrounds of the described variety.

115

Discussion and Collaboration

116

TALK: SPLENDOR: Narrow-gap Quantum Materials for Light Dark Matter

Corresponding Author: yfkahn@illinois.edu

117

TALK: SPLENDOR: narrow-gap quantum materials for light dark matter**Corresponding Author:** yfkahn@illinois.edu

Author: Yonatan Kahn

Low-mass direct detection experiments based on conventional semiconductors such as silicon and germanium are achieving impressive sensitivity for dark matter masses above 1 MeV. However, to probe dark matter-electron scattering via electronic excitations for even lighter dark matter masses, more exotic materials with sub-eV band gaps are required. I will give an overview of the SPLENDOR program, which aims to develop a prototype detector for keV-MeV dark matter based on narrow-gap quantum materials. This research program involves close interdisciplinary collaboration with condensed matter theorists and experimentalists, in all stages from detector material synthesis to determining the experimental sensitivity via the density response function.

118

Shuttle to Middle School for Discussion and Collaboration

119

TALK: TBD

. / 120

Discussion and Collaboration

121

TALK: Dynamical Generation of the Baryon Asymmetry from a Scale Hierarchy**Corresponding Author:** jhchang@fnal.gov

Authors: Jae Hyeok Chang, Kwang Sik Jeong, Chang Hyeon Lee, Chang Sub Shin

We propose a novel baryogenesis scenario where the baryon asymmetry originates directly from a hierarchy between two fundamental mass scales: the electroweak scale and the Planck scale. Our model is based on the neutrino-portal Affleck-Dine (AD) mechanism, which generates the asymmetry of the AD sector during the radiation-dominated era and subsequently transfers it to the baryon number before the electroweak phase transition. The observed baryon asymmetry is then a natural outcome of this scenario. The model is testable as it predicts the existence of a Majoron with a keV mass and an electroweak scale decay constant. The impact of the relic Majoron on ΔN_{eff} can be measured through near-future CMB observations.

122

TALK: Gauged global strings

Corresponding Author: weixue@ufl.edu

Author: Wei Xue

I will present the string solutions and cosmological implications of the gauge $U(1)_Z \times$ global $U(1)_{PQ}$ model. With two hierarchical symmetry-breaking scales, the model exhibits three distinct string solutions: a conventional global string, a global string with a heavy core, and a gauge string as a bound state of the two global strings. This model reveals rich phenomenological implications in cosmology. When incorporating this model with the QCD axion framework, the heavy-core global strings emit more axion particles due to their large tension. This radiation significantly enhances the QCD axion dark matter abundance, thereby opening up the QCD axion mass window. Furthermore, in contrast to conventional gauge strings, the gauge strings in this model exhibit a distinctive behavior, radiating axions.

123

Discussion, Collaboration, and Lunch

124

TALK: Cosmic Stability of Dark Matter from Pauli Blocking

Corresponding Author: batell@pitt.edu

Why does dark matter (DM) live longer than the age of the Universe? Here we study a novel sub-eV scalar DM candidate whose stability is due to the Pauli exclusion of its fermionic decay products. We analyze the stability of the DM condensate against decays, scatterings (i.e., evaporation), and parametric resonance, delineating the viable parameter regions in which DM is cosmologically stable. In a minimal scenario in which the scalar DM decays to a pair of new exotic fermions, we find that scattering can populate an interacting thermal dark sector component to energies far above the DM mass. This self-interacting dark radiation may potentially alleviate the Hubble tensions. Furthermore, our scenario can be probed through precise measurements of the halo mass function or the masses of dwarf spheroidal galaxies since scattering prevents the DM from becoming too dense. On the other hand, if the lightest neutrino stabilizes the DM, the cosmic neutrino background can be significantly altered from the standard cosmology and thus be probed in the future by cosmic neutrino background detection experiments.

125

Discussion and Collaboration

126

TALK: BSM Physics with Gravitational Wave Detectors

Corresponding Author: kuver.sinha@ou.edu

Author: Kuver Sinha

Future gravitational wave detectors probing the mHz - nHz frequency range will provide a unique opportunity for BSM physicists to study new physics. Neutron star and white dwarf mergers can serve as axion probes, while extreme mass ratio inspirals can constrain dark forces. Gravitational wave detectors will also probe early first order phase transitions. I will discuss some ongoing work and future ideas in these directions.

128

Discussion, Collaboration, and Lunch

129

TALK: Cosmological Stasis and Its Realization from Dynamical Scalars

Corresponding Author: thomasbd@lafayette.edu

Author: Brooks Thomas

Co-authors: Keith R. Dienes, Lucien Heurtier, Fei Huang, Timothy M. P. Tait

It has recently been realized that many extensions of the Standard Model give rise to cosmological histories exhibiting extended epochs of cosmological stasis —epochs wherein the abundances of multiple energy components (such as matter, radiation, or vacuum energy) remain effectively constant despite cosmological expansion. The emergence of a stasis epoch is not a consequence of fine-tuning in cosmologies of this sort; rather, stasis turns out to be a global attractor toward which the universe naturally evolves for a broad range of initial conditions. In this talk, I shall review the general conditions under which stasis emerges in such scenarios and explore some of its potential implications for cosmological observables such as the matter power spectrum and the stochastic gravitational-wave background. I shall also discuss a particular realization of stasis involving a collection of scalar fields, each of which dynamically transitions from a period of slow roll to a period of rapid oscillation around its potential minimum as the universe expands. As I shall demonstrate, not only does cosmological stasis arise in such scenarios, but the system of dynamical scalars also exhibits novel features not seen in previous realizations of stasis, including a tracking behavior wherein the effective equation of state for the universe as a whole evolves toward the equation of state of this energy component. The emergence of such tracking behavior has potential model-building implications in the context of dark-energy and cosmic-inflation scenarios.

130

TALK: Updates on the Migdal effect and hydrogen doping for dark matter detection

Corresponding Author: jayden.newstead@unimelb.edu.au

Author: Jayden Newstead

An ongoing challenge in dark matter direct detection is to improve the sensitivity to light dark mat-

ter in the MeV–GeV mass range. One proposal is to dope a liquid noble-element direct-detection experiment with a lighter element such as hydrogen. This has the advantage of enabling larger recoil energies compared to scattering on a heavy target, while leveraging existing detector technologies. Direct-detection experiments can also extend their reach to lower masses by exploiting the Migdal effect, where a nuclear recoil leads to electronic ionization or excitation. In this work, we combine these ideas to study the sensitivity of a hydrogen-doped LZ experiment (HydroX) and a future large-scale experiment such as XLZD. We find that HydroX could have sensitivity to dark matter masses below 10 MeV for both spin-independent and spin-dependent scattering, with XLZD extending that reach to lower cross sections. Notably, this technique substantially enhances the sensitivity of direct detection to spin-dependent proton scattering, well beyond the reach of any current experiments.

131

Discussion and Collaboration

132

TALK: Dark Matter Raining on DUNE and Other Large Volume Detectors

Corresponding Author: joshua.berger@colostate.edu

Authors: Javier Acevedo, Joshua Berger, Peter Denton

Direct detection is a powerful means of searching for particle physics evidence of dark matter (DM) heavier than about a GeV with volume, low-threshold detectors.

In many scenarios, some fraction of the DM may be boosted to large velocities enhancing and generally modifying possible detection signatures. We investigate the scenario where 100% of the DM may be boosted at the Earth due to new attractive long-range forces. This opens up two main improvements in detection capabilities: 1) the detection signatures are stronger opening up large-volume neutrino detectors, such as DUNE, Super-K, Hyper-K, and JUNO, as possible DM detectors, and 2) the large boost allows for detectable signatures of sub-GeV DM. At lower boosts, a modified, higher-than-usual energy signal could be accessible at direct detection experiments such as LZ. In addition, the model leads to a significant anisotropy in the signal with the DM flowing dominantly vertically at the Earth's surface instead of the typical approximately isotropic DM signal. We develop the theory behind this model and also calculate realistic constraints using a detailed GENIE simulation of the signal inside detectors.

133

TALK: Gravitational Waves from Cosmic Superstrings and Gauge Strings

Corresponding Author: gnaoiypb@duck.com

Author: Danny Marfatia

We perform a phenomenological comparison of the gravitational wave spectrum expected from cosmic gauge string networks and superstring networks comprised of multiple string types.

134

Discussion, Collaboration, and Lunch

135

TALK: Introduction to DAMSA, A Novel Dark Matter Search Experiment at an Accelerator

Corresponding Author: jaehoonyu@uta.edu

Dark matter is thought to make up 25% of the universe. Dark sector particles (DSP) do not interact through the known forces but could be weakly coupled to Standard Model particles through a portal or a mediator. Many searches for dark matter/dark sector particles at an accelerator thus far seem to face a ceiling that the sensitivity reach is greatly limited, beyond statistical effects. DAMSA is an extremely short baseline experiment that proposes to break through this limit, taking advantage of high beam powers available at various accelerator facilities around the world, including the PIP-II Linac under construction, an essential element in providing the necessary high flux proton beams to DUNE at Fermilab. In this talk, I will describe the DAMSA (Dump produced Aboriginal Matter Search at an Accelerator) experiment. I will also discuss current status and plan for DAMSA and its expected sensitivity reach in the search of the Axion-Like Particle as an example physics case.

136

Discussion and Collaboration

137

TALK: Dark Matter in the Time of Gravitational Waves

Corresponding Author: tao.xu@ou.edu

The observation of gravitational waves opens a new window for exploring astrophysics and cosmology. These messengers enable the concurrent measurement of their amplitudes and phases, facilitating a precise analysis of gravitational wave production and propagation. In this talk, I will demonstrate how gravitational waves can be utilized to study the properties of dark matter. Specifically, I will use wave dark matter as an example to show that gravitational waveforms, along with further multi-messenger observations involving photon signals, reveal distinctive features. These features can be probed with the ongoing LIGO and upcoming LISA missions.

138

TALK: Probing self-interacting sterile neutrino dark matter with the diffuse supernova neutrino background

Corresponding Author: asuliga@berkeley.edu

Author: Anna Suliga

Co-authors: Baha Balantekin, George Fuller, Anupam Ray

The neutrinos in the diffuse supernova neutrino background (DSNB) travel over cosmological distances and this provides them with an excellent opportunity to interact with dark relics. We show that a cosmologically-significant relic population of keV-mass sterile neutrinos with strong self-interactions could imprint their presence in the DSNB. The signatures of the self-interactions would be “dips” in the otherwise smooth DSNB spectrum. Upcoming large-scale neutrino detectors, for example Hyper-Kamiokande, have a good chance of detecting the DSNB and these dips. If no dips are detected, this method serves as an independent constraint on the sterile neutrino self-interaction strength and mixing with active neutrinos. We show that relic sterile neutrino parameters that evade X-ray and structure bounds may nevertheless be testable by future detectors like TRISTAN, but may also produce dips in the DSNB which could be detectable. Such a detection would suggest the existence of a cosmologically-significant, strongly self-interacting sterile neutrino background, likely embedded in a richer dark sector.

139

Discussion, Collaboration, and Lunch

140

TALK: Introduction to DAMSA, A Novel Dark Matter Search Experiment at an Accelerator

Corresponding Author: jaehoonyu@uta.edu

141

Discussion and Collaboration

142

TALK: Journey with Sourced Millicharged Particles and Dark Photons

Corresponding Author: zliuphys@umn.edu

Author: Zhen Liu

Hidden symmetries are widely prevalent across numerous new physics models and have garnered significant attention. These symmetries give rise to intriguing testable phenomena, such as “dark photons” as the force mediators and “millicharged” particles representing the matter content. A diverse range of physics experiments and search strategies have been developed to investigate these phenomena. In this talk, I will discuss my recent work in exploring these elusive particles, covering both theoretical foundations and experimental results. I will focus on the unique aspects of controlled search approaches, where we actively source the fields under investigation and discuss the future prospects in this exciting research area.

143

TALK: Asymptotically Free E_6 GUT and the Generation of Neutrino Mass

Corresponding Author: vasja.susic@lnf.infn.it

Author: Vasja Susič

Co-authors: K.S Babu, Borut Bajc

Grand Unified Theories (GUTs) represent an attractive possibility of physics beyond the Standard Model, with implications for both proton decay and neutrinos. The most studied examples are based on unified groups $SU(5)$ and $SO(10)$, while the exceptional case remains less understood. In an asymptotically free GUT, model building is essentially limited to the use of irreducible representations of dimension 27 and 78 (the fundamental and the adjoint). I will present ongoing work in non-supersymmetric models regarding the determination of the necessary ingredients for both GUT symmetry breaking and a realistic Yukawa sector. Special attention will be given to the peculiarities associated with neutrino mass generation in this type of models.

144

Discussion, Collaboration, and Lunch

145

TALK: TALK: Gravitational Wave Symphony from Oscillating Spectator Scalar Fields

Corresponding Author: yanou.cui@ucr.edu

Spectator scalar fields can be generically present in the early Universe, and are potentially dark matter candidates. We investigate the prospect of such scalars loop as a generic source of stochastic gravitational wave background (SGWB) due to parametric resonance during their oscillation phase. By systematically analyzing benchmark models through lattice simulations and considering a wide range of parameters, we demonstrate that such a scenario can lead to detectable signals in GW detectors over a broad frequency range and potentially address the recent findings by Pulsar Timing Array experiments. Furthermore, we show that these models naturally yield viable ultra-light, wave-like dark matter candidates and/or dark radiation detectable by CMB observatories. We also explore the potential for realizing low-scale baryogenesis from such an oscillating scalar system and the resultant detectable SGWB signals. This study highlights an intriguing connection between the stochastic gravitational wave background (SGWB) and terrestrial probes for low-energy new particle physics.

146

Discussion and Collaboration

147

TALK: Probing Baryon Number Violation in Neutron Stars**Corresponding Author:** rouzbeh@unm.edu

Author: Rouzbeh Allaverdi

I start by presenting minimal extensions of the Standard Model that accommodate low-scale baryogenesis. These models involve fermion singlets with baryon-number-violating couplings to quarks. Then I briefly discuss the consequences of these models for low energy and high energy experiments. Finally, I mention how neutron stars can be used to tightly constrain such interactions for GeV scale fermions.

148

TALK: TBD (open slot available)

149

Discussion, Collaboration, and Lunch

150

TALK: TBD (open slot available)

151

Discussion and Collaboration

152

TALK: Does the Sun Have a Dark Disk?**Corresponding Author:** m.zakeri@uky.edu

The Sun's oblateness has been measured using optical observations. Its gravitational quadrupole moment has been deduced through helioseismology and measurements of its gravitational effects on Mercury's orbit. The distribution of mass within Mercury's orbit would only impact the orbital determination, suggesting that discrepancies among various types of assessments may indicate the possible existence of a non-luminous mass. For the first time, we have developed a method to combine these differing measurements to yield new, highly sensitive constraints on the mass distribution within Mercury's orbit. In this talk, we will show that the most precise measurements indicate the existence of a non-luminous disk within Mercury's orbit, with a mass significantly heavier than the

modeled mass for the circumsolar dust ring observed by the Solar TERrestrial RELations Observatory (STEREO) mission. This suggests a substantial dark matter contribution. Furthermore, the long-standing inconsistency between the element abundances determined from the spectroscopy of the Sun's surface, and those inferred from its interior through helioseismology, can be reconciled if the Sun formed within a protoplanetary disk. We will discuss how our findings limit the presence of a dark disk or a spherical halo near the Sun and highlight the potential of future orbital measurements of Mercury and near-Sun asteroids to refine these constraints further.

153

TALK: TBD (open slot available)

154

Discussion, Collaboration, and Lunch

155

Independence Day Festivities in Lead

156

TBD (open slot available)

157

TALK: Neutrino Interactions**Corresponding Author:** vpandey@fnal.gov

158

Discussion, Collaboration, and Lunch

159

TALK: Title TBD (Speaker: Volodymyr Takhistov)

Corresponding Author: vtakhist@post.kek.jp

160

Discussion and Collaboration

161

The Institute for Underground Science at SURF Update

162

TALK: CP - Violation with Neutrino Disappearance

Corresponding Author: peterbd1@gmail.com

Author: Peter Denton

The best way to probe CP violation in the lepton sector is with long-baseline accelerator neutrino experiments in the appearance mode: the appearance of in predominantly beams. Here we show that it is possible to discover CP violation with disappearance experiments only, by combining JUNO for electron neutrinos and DUNE or Hyper-Kamiokande for muon neutrinos. While the maximum sensitivity to discover CP is quite modest (with 6 years of JUNO and 13 years of DUNE), some values of may be disfavored by depending on the true value of

163

Discussion, Collaboration, and Lunch

164

TALK: A 17 MeV pseudoscalar and the MiniBooNE, LSND and ATOMKI anomalies

In the absence of any new physics signals at the Large Hadron Collider (LHC), anomalous results at low energy experiments have become the subject of increased attention and scrutiny. We focus on three such results from the LSND, MiniBooNE (MB), and ATOMKI experiments. A 17 MeV pseudoscalar mediator can account for the excess events seen in 8Be and 4He pair creation transitions in ATOMKI. We incorporate this mediator in a gauge invariant extension of the Standard Model (SM) with a second Higgs doublet and three singlet (seesaw) neutrinos (N_i , $i = 1, 2, 3$). $N_{1,2}$ participate in an interaction in MB and LSND which, with a ϕ as mediator, leads to the production of e^+e^- pairs. The N_i also lead to mass-squared differences for SM neutrinos in agreement with global oscillation

data. We first show that such a model offers a clean and natural joint solution to the MB and LSND excesses. We then examine the possibility of a common solution to all three anomalies. Using the values of the couplings to the quarks and electrons which are required to explain pair creation nuclear transition data for ^8Be and ^4He in ATOMKI, we show that these values lead to excellent fits for MB and LSND data as well, allowing for a common solution. We discuss the constraints and future tests of our proposal.

165

Discussion and Collaboration

166

TALK: Charged-current (anti)neutrino-nucleon Scattering and QED Nuclear Medium

Corresponding Author: tomalak@lanl.gov

Author: Oleksandr Tomalak

Charged-current quasielastic neutrino scattering is the signal process in neutrino oscillation experiments and requires precise theoretical prediction for the analysis of modern and future experimental data, starting with the nucleon axial-vector form factor. In this talk, I compare a new MINERvA measurement of this form factor with lattice-QCD calculations and deuterium bubble-chamber data, provide uncertainty projections for future extractions, present recent calculations of radiative corrections to charged-current processes, and investigate the potential of neutrino scattering data on constraining nucleon- and quark-level interactions beyond the Standard Model. The exchange of photons with nuclear medium modifies (anti)neutrino and electron scattering cross sections. We study the distortion of (anti)neutrino-nucleon and charged lepton-nucleon cross sections, medium-induced bremsstrahlung, and estimate the QED-medium effects on the final-state kinematics and scattering cross sections. We find new permille-to-percent level effects, which were never accounted for in either (anti)neutrino-nucleon or electron-nucleon scattering. We quantitatively compute the effects of Glauber photon-mediated multiple re-scattering within the nuclear medium and find that the relativistic charged lepton acquires a momentum of order 10 MeV transverse to its direction of propagation inside the nucleus. This broadening sizably deflects expected electron and muon tracks and suppresses scattering cross sections. Precise extraction of the nucleon and nuclear structure by electron and muon probes should, thus, take the QED nuclear medium angular redistribution of particles into account.

167

TALK: Constraining Non-standard Neutrino Interactions with Neutral Current Events of Long-Baseline Oscillation Experiments

Corresponding Author: julia.gehrlein@colostate.edu

Author: Julia Gehrlein

Neutrino non-standard interactions (NSI) provide an attractive framework to study new physics in the neutrino sector in scattering experiments and in oscillation experiments where NSI lead to new matter effects. At long-baseline oscillation experiments, so far, constraints on NSI have been

derived using charged current events. In this manuscript we explore a new channel, namely neutral-current events at long-baseline experiments to constrain vector and axial-vector NSI. We introduce a framework to parametrize the effect of NSI on the cross section and then focus, as an example, on the NOvA experiment to derive constraints on NSI using real data.

168

Discussion, Collaboration, and Lunch

169

TALK: A SURF Low Background Module (SLoMo)

Corresponding Author: juergen.reichenbacher@sdsmt.edu

With radiopurity controls and targeted design modifications a kton-scale liquid argon time projection chamber similar to DUNE could be used for enhanced low energy physics searches. This includes improved sensitivity to supernova and solar neutrinos, and other rare event searches, such as WIMP dark matter, while simultaneously serving long-baseline neutrino physics. This talk will present simulation studies to evaluate physics sensitivities of such a module. It will also discuss R&D to develop large-scale radiopurity controls necessary to construct such a detector.

170

Discussion and Collaboration

171

Shuttle to Middle School

172

Shuttle to Sanford Lab Visitor Center

173

TALK: Probing Heavy Neutral Lepton on Muon Collider and Beam-dump Experiments

Corresponding Author: lyu00145@umn.edu

Author: Kunfeng Lyu

Broad classes of solutions to the neutrino puzzles can be best tested by seeking the partners of SM light neutrinos, dubbed as heavy neutral leptons (HNLs). It can be parameterized by the HNL mass m_N and the mixing angle U_1 with the SM neutrino. In this talk, I will mainly discuss about probing two parameter space regime. One is for HNL mass larger than $O(100)$ GeV which can be tested at future high energy muon collider. Alternatively, if the HNL is lighter than the muon, it can be produced by the decays of muons and pions. The LSND experiment and the future beam dump experiment such as PIP2-BD can be exploited to improve the sensitivity.

174

Discussion, Collaboration, and Lunch

175

Shuttle to Ethnobotanical Garden

176

Guest Speaker at Ethnobotanical Garden - Rochelle Zens

Corresponding Author: rzens@sanfordlab.org

177

Shuttle to Middle School

178

TALK: Neutrino and light dark matter physics with directional detectors

Corresponding Author: diego.aristizabal@usm.cl

Author: Diego Aristizabal

In this talk I will review a few aspects of Coherent Elastic neutrino-Nucleus Scattering (CEvNS) and light dark matter (LDM) production in meson decays. In particular, I will focus on potential measurements using neutrino beams at FNAL. I will show that, combined with directional detectors, these beams offer an avenue for CEvNS measurements as well as for LDM detection. Such measurements will provide complementary information to that coming from other CEvNS facilities as well as from other non-directional detectors looking for LDM.

179

Discussion and Collaboration

180

TALK: Title TBD (Speaker: Garv Chauhan)

Corresponding Author: garv.chauhan@gmail.com

181

TALK: The Scotos Side of Neutrinos

Corresponding Author: schakdar@holycross.edu

Authors: Kaladi Babu, Shreyashi Chakdar, Vishnu Padmanabhan Kovilakam

We carry out a systematic investigation for minimal Scotogenic models based on a dark gauge symmetry, in which the neutrino masses are induced at the one-loop level and include a chiral dark matter (DM) candidate. Assuming this gauge symmetry is broken by only one Higgs singlet scalar that also generates masses to all dark fermions, we analyze the stability of the DM candidate which is ensured by a residual symmetry of gauge symmetry. There can be different DM scenarios explored in this framework and we investigate the associated scalar and fermionic DM phenomenology of one of the minimal models.

182

Discussion, Collaboration, and Lunch

183

TALK: Does the Sun have a dark disk?

Corresponding Author: gardner@pa.uky.edu

Authors: Gustavo Alves, Susan Gardner, Pedro Machado, Mohammadreza Zakeri

The nature of dark matter is unknown, as are its total mass and distribution within the Solar system, though constraints from measurements of planetary orbits exist. I discuss how differing measurements of the Sun's gravitational quadrupole moment can combine to yield new and highly sensitive constraints on the mass distribution within Mercury's orbit. The best determinations point with high confidence to the existence of a non-luminous disk coplanar with Mercury's orbit, and the mass estimates associated with known matter, although uncertain, point to a significant dark-matter contribution, for which macroscopic dark-matter models may suit best.

184

Discussion and Collaboration

185

Coffee/Social with Students

186

TALK: Direct Collapse Black Holes from Dark Matter Annihilation

Corresponding Author: flip.tanedo@ucr.edu

187

TALK: Quasi (pseudo)-Dirac Neutrinos

Corresponding Author: sheng.fong@ufabc.edu.br

Author: Chee Sheng Fong

Here we will consider quasi (pseudo)-Dirac neutrinos

188

Discussion and Collaboration

189

TALK: Neutrinos From the Sun Can Discover Dark Matter-electron Scattering

Corresponding Author: ranjanlaha@iisc.ac.in

Author: Ranjan Laha

We probe dark matter-electron scattering using high-energy neutrino observations from the Sun. Dark matter (DM) interacting with electrons can get captured inside the Sun. These captured DM may annihilate to produce different Standard Model (SM) particles. Neutrinos produced from these SM states can be observed in IceCube and DeepCore. Although there is no excess of neutrinos from the Solar direction, we find that the current data-sets of IceCube and DeepCore set the strongest constraint on DM-electron scattering cross section in the DM mass range 10 GeV to 10

GeV. Our work implies that future observations of the Sun by neutrino telescopes have the potential to discover DM-electron interaction.

190

TALK: E6 Unification: Intermediate Symmetries, Fermion Masses and Proton Decay

Corresponding Author: borut.bajc@ijs.si

Authors: Kaladi Babu, Borut Bajc, Vasja Susič

I will present a non-supersymmetric E6 GUT with the scalar sector consisting of $650 + 351' + 27$ dimensional representations. The intermediate symmetries which turn out to be realistic under the extended survival hypothesis (with minimal fine-tuning) are trinification symmetry $SU(3)_C \times SU(3)_L \times SU(3)_R$ with either LR or CR parity, and $SU(6)_{CR} \times SU(2)_L$. This means among others that they can reproduce correctly the light charged fermion and neutrino masses as well as the CKM and PMNS mixing matrices. Although the successful cases give a large range for proton lifetime estimates, all of them include regions consistent with current experimental bounds and within reach of forthcoming experiments.

191

Discussion, Collaboration, and Lunch

192

TALK: TBD (open slot available)

193

Discussion and Collaboration

194

TALK: New Directions for ALP Searches Combining Nuclear Reactors and Haloscopes

Corresponding Author: vedran.brdar@okstate.edu

Author: Vedran Brdar

I will discuss a recently proposed novel experimental setup for axion-like particle (ALP) searches.

Nuclear reactors produce a copious number of photons, a fraction of which could convert into ALPs via the Primakoff process in the reactor core. The generated flux of ALPs leaves the nuclear power plant, and its passage through a region with a strong magnetic field results in efficient conversion to photons, which can be detected. Such a magnetic field is the key component of axion haloscope experiments. I will discuss existing setups featuring an adjacent nuclear reactor and axion haloscope and I will demonstrate that the obtained sensitivity projections complement constraints from existing laboratory experiments, e.g., light-shining-through-walls.

195

TALK: Decoding the Flavor Evolution of Supernova Neutrinos: Shedding Light on Neutrino Electromagnetic Properties and Supernova Dynamics

Corresponding Author: yago.porto@ufabc.edu.br

Author: Yago Porto

Two of the most important questions in neutrino astrophysics are: A) How can the extreme supernova environments, through which neutrinos propagate, be used to shed light on the new physics behind neutrino masses and mixing? B) What is the flavor evolution of supernova neutrinos in the latest stages of core collapse? In this talk, I will summarize our recent developments in these directions, emphasizing how the unique magnetic fields of supernova progenitors can modify neutrino evolution in the presence of both Dirac and Majorana neutrino magnetic moments during the first few tens of milliseconds of neutrino emission. For the rest of the neutrino emission (roughly 10 seconds), I will describe how standard matter effects can define the main features of the neutrino signal in terrestrial detectors despite the exotic neutrino-neutrino refraction that dominates in the supernova core.

196

Discussion, Collaboration, and Lunch

197

TALK: TBD (open slot available)

198

Discussion and Collaboration

199

TALK: Probing SUSY at Gravitational Wave Observations

Corresponding Author: shaikh.saad@okstate.edu

Author: Shaikh Saad

Under the assumption that the recent pulsar timing array evidence for a stochastic gravitational wave (GW) background at nanohertz frequencies is generated by metastable cosmic strings, we analyze the potential of present and future GW observatories for probing the change of particle degrees of freedom caused, e.g., by a supersymmetric (SUSY) extension of the Standard Model (SM). We find that signs of the characteristic doubling of degrees of freedom predicted by SUSY could be detected at Einstein Telescope and Cosmic Explorer even if the masses of the SUSY partner particles are as high as about TeV, far above the reach of any currently envisioned particle collider. We also discuss the detection prospects for the case that some entropy production, e.g. from a late decaying modulus field inducing a temporary matter domination phase in the evolution of the universe, somewhat dilutes the GW spectrum, delaying discovery of the stochastic GW background at LIGO-Virgo-KAGRA. In our analysis we focus on SUSY, but any theory beyond the SM predicting a significant increase of particle degrees of freedom could be probed this way.

200

TALK: Embedding the Universal See-Saw Mechanism in Pati-Salam Model

Corresponding Author: sumit.biswas@okstate.edu

Authors: Kaladi S. Babu, Sumit Biswas

Here we explore the integration of universal see-saw mechanism within the parity-symmetric Pati-Salam framework to address fermion mass generation. Incorporating vector-like fermions and a simple Higgs sector helps realize the hierarchical mass for all generations in this unified quark-lepton model. The neutrino sector seems pretty intriguing with the lightest neutrino acquiring mass through one-loop radiative corrections. With softly broken parity, we also see the possibility of addressing the strong CP problem in this setup.

201

Discussion, Collaboration, and Lunch

202

TALK: eV Sterile Neutrino Paradox

Corresponding Author: sudip.jana@okstate.edu

Authors: Sudip Jana, Lucas Puetter, Alexei Yu. Smirnov

Light sterile neutrinos are frequently featured in various theories, particularly to explain short-baseline anomalies or as candidates for dark matter. These scenarios necessitate non-zero mixing with active neutrinos, often disregarding the contributions to the active neutrino mass matrix. We demonstrate that this mixing induces contributions to the active neutrino mass matrix, thereby affecting neutrino oscillation results. We conduct a rigorous analysis and present stringent bounds on these induced masses for each entry of a active neutrino mass matrix. Furthermore, we examine whether the induced matrix can elucidate the distinctive pattern of lepton mixing and the neutrino mass spectrum, while discussing the various phenomenological implications.

203

Discussion and Collaboration

204

Shuttle Pickup at Hampton Inn, travel to Sanford Lab Homestake Visitor Center

The workshop will be held at the visitor center all day today.

205

Shuttle Pickup at Middle School, Travel to E&O Building

206

SURF Overview by Mike Headley, Executive Director, South Dakota Science and Technology Authority

207

SURF Underground Tour Safety, Waivers, Lunch at 11:00 w/ Cultural Awareness Committee, PPE and Shuttle to Ross

Corresponding Author: jmolina@sanfordlab.org

Only the CETUP* participants that are going on the underground tour will attend. Other participants not going on the tour will be able to stay at the Middle School and continue to work and carry on discussions.

208

Shuttle, PPE, Return to Hampton Inn

209

TALK: TBD (open slot available)

210

TALK: Skimming tau neutrinos and optical Cherenkov signals**Corresponding Author:** mary-hall-reno@uiowa.edu

Author: Mary Hall Reno

Optical Cherenkov signals from up-going air showers show promise for detection of skimming tau neutrinos. We review the theoretical inputs to evaluating the sensitivities to transient neutrino sources of orbital and sub-orbital Cherenkov telescopes.

211

Discussion, Collaboration, and Lunch

212

TALK: TBD (open slot available)

213

Discussion and Collaboration

214

Road Scholar Group - CETUP* Guest Speaker (Mary Hall Reno)**Corresponding Author:** mary-hall-reno@uiowa.edu

CETUP 2024 / 215

Dynamical Generation of the Baryon Asymmetry from a Scale Hierarchy**Authors:** Chang Hyeon Lee¹; Chang Sub Shin¹; Jae Hyeok Chang²; Kwang Sik Jeong³¹ *Chungnam Natl. U.*² *Fermilab and UIC*³ *Pusan Natl. U.*

Corresponding Author: jhchang@fnal.gov

We propose a novel baryogenesis scenario where the baryon asymmetry originates directly from a hierarchy between two fundamental mass scales: the electroweak scale and the Planck scale. Our model is based on the neutrino-portal Affleck-Dine (AD) mechanism, which generates the asymmetry of the AD sector during the radiation-dominated era and subsequently transfers it to the baryon number before the electroweak phase transition. The observed baryon asymmetry is then a natural outcome of this scenario. The model is testable as it predicts the existence of a Majoron with a keV mass and an electroweak scale decay constant. The impact of the relic Majoron on ΔN_{eff} can be measured through near-future CMB observations.

216

Discussion and Collaboration

CETUP 2024 / 217

The Power of the Dark Sink

Author: Robert McGehee¹

¹ *University of Minnesota*

Corresponding Author: rmcgehee@umn.edu

We describe a simple dark sector structure which, if present, has implications for the direct detection of dark matter (DM): *the Dark Sink*. A Dark Sink transports energy density from the DM into light dark-sector states that do not appreciably contribute to the DM density. As an example, we consider a light, neutral fermion ψ which interacts solely with DM χ via the exchange of a heavy scalar Φ . We illustrate the impact of a Dark Sink by adding one to a DM freeze-in model in which χ couples to a light dark photon γ' which kinetically mixes with the Standard Model (SM) photon. This freeze-in model (absent the sink) is itself a benchmark for ongoing experiments. In some cases, the literature for this benchmark has contained errors; we correct the predictions and provide them as a public code. We then analyze how the Dark Sink modifies this benchmark, solving coupled Boltzmann equations for the dark-sector energy density and DM yield. We check the contribution of the Dark Sink ψ 's to dark radiation; consistency with existing data limits the maximum attainable cross section. For DM with a mass between MeV – $\mathcal{O}(10 \text{ GeV})$, adding the Dark Sink can increase predictions for the direct detection cross section all the way up to the current limits.

CETUP 2024 / 218

Analytic Results on Dark Matter Velocity Distributions

Author: Jason Kumar¹

¹ *University of Hawaii*

Corresponding Author: jkumar@hawaii.edu

We review recent work on determining dark matter velocity distributions from analytic methods of classical mechanics, and the potential impact of these results on dark matter indirect detection strategies. In particular, we discuss velocity-dependent dark matter annihilation in subhalos and in the Galactic Center. We compare the results of these analytic methods to those obtained in large N numerical simulations, and discuss future work.

CETUP 2024 / 219

Probing SUSY at Gravitational Wave Observatories

Author: Shaikh Saad^{None}

Corresponding Author: shaikh.saad@okstate.edu

Under the assumption that the recent pulsar timing array evidence for a stochastic gravitational wave (GW) background at nanohertz frequencies is generated by metastable cosmic strings, we analyze the potential of present and future GW observatories for probing the change of particle degrees of freedom caused, e.g., by a supersymmetric (SUSY) extension of the Standard Model (SM). We find that signs of the characteristic doubling of degrees of freedom predicted by SUSY could be detected at Einstein Telescope and Cosmic Explorer even if the masses of the SUSY partner particles are as high as about 10^4 TeV, far above the reach of any currently envisioned particle collider. We also discuss the detection prospects for the case that some entropy production, e.g. from a late decaying modulus field inducing a temporary matter domination phase in the evolution of the universe, somewhat dilutes the GW spectrum, delaying discovery of the stochastic GW background at LIGO-Virgo-KAGRA. In our analysis we focus on SUSY, but any theory beyond the SM predicting a significant increase of particle degrees of freedom could be probed this way.

220

-TBD-

Author: Garv Chauhan¹

¹ *Virginia Tech*

Corresponding Author: garv.chauhan@gmail.com

N/A

221

Gravitational waves from cosmic superstrings and gauge strings

Author: Danny Marfatia^{None}

Corresponding Author: gnaoiypb@duck.com

We perform a phenomenological comparison of the gravitational wave spectrum expected from cosmic gauge string networks and superstring networks comprised of multiple string types.

222

Journey with Sourced Millicharged Particles and Dark Photons

Author: Zhen Liu¹

¹ *University of Minnesota*

Corresponding Author: zliuphys@umn.edu

Hidden $U(1)$ symmetries are widely prevalent across numerous new physics models and have garnered significant attention. These symmetries give rise to intriguing testable phenomena, such as “dark photons” as the force mediators and “millicharged” particles representing the matter content. A diverse range of physics experiments and search strategies have been developed to investigate these phenomena. In this talk, I will discuss my recent work in exploring these elusive particles, covering both theoretical foundations and experimental results. I will focus on the unique aspects of controlled search approaches, where we actively source the fields under investigation and discuss the future prospects in this exciting research area.

223

gauged global strings

Author: Wei Xue¹

¹ *University of Florida*

Corresponding Author: weixue@ufl.edu

I will present the string solutions and cosmological implications of the gauge $U(1)Z \times \text{global } U(1)PQ$ model. With two hierarchical symmetry-breaking scales, the model exhibits three distinct string solutions: a conventional global string, a global string with a heavy core, and a gauge string as a bound state of the two global strings. This model reveals rich phenomenological implications in cosmology. When incorporating this model with the QCD axion framework, the heavy-core global strings emit more axion particles due to their large tension. This radiation significantly enhances the QCD axion dark matter abundance, thereby opening up the QCD axion mass window. Furthermore, in contrast to conventional gauge strings, the gauge strings in this model exhibit a distinctive behavior, radiating axions.

224

Probing Baryon Number Violation in Neutron Stars

Author: Rouzbeh Allaverdi¹

¹ *University of New Mexico*

Corresponding Author: rouzbeh@unm.edu

I start by presenting minimal extensions of the Standard Model that accommodate low-scale baryogenesis. These models involve fermion singlets with baryon-number-violating couplings to quarks. Then I briefly discuss the consequences of these models for low energy and high energy experiments. Finally, I mention how neutron stars can be used to tightly constrain such interactions for GeV scale fermions.

225

Welcome by Mike Headley, Executive Director, South Dakota Science and Technology Authority/SURF

226

Shuttle to Hampton Inn

227

Modular Flavor Symmetry and Neutrino Oscillations

Author: Zurab Tavartkiladze¹

¹ *Ilia State University*

Corresponding Author: zurab.tavartkiladze@iliauni.edu.ge

Extension of the SM with modular flavor symmetry will be discussed. We show that by specific assignment of the transformation properties to the fermion states, very economical and predictive lepton sector can be constructed. Details of the neutrino sector and also robustness of the obtained results will be discussed.

228

SPLENDOR: narrow-gap quantum materials for light dark matter

Author: Yonatan Kahn¹

¹ *University of Illinois Urbana-Champaign*

Corresponding Author: yfkahn@illinois.edu

Low-mass direct detection experiments based on conventional semiconductors such as silicon and germanium are achieving impressive sensitivity for dark matter masses above 1 MeV. However, to probe dark matter-electron scattering via electronic excitations for even lighter dark matter masses, more exotic materials with sub-eV band gaps are required. I will give an overview of the SPLENDOR program, which aims to develop a prototype detector for keV-MeV dark matter based on narrow-gap quantum materials. This research program involves close interdisciplinary collaboration with condensed matter theorists and experimentalists, in all stages from detector material synthesis to determining the experimental sensitivity via the density response function.

229

Updates on the Migdal effect and hydrogen doping for dark matter detection

Author: Jayden Newstead¹

¹ *University of Melbourne*

Corresponding Author: jayden.newstead@unimelb.edu.au

An ongoing challenge in dark matter direct detection is to improve the sensitivity to light dark matter in the MeV–GeV mass range. One proposal is to dope a liquid noble-element direct-detection experiment with a lighter element such as hydrogen. This has the advantage of enabling larger

recoil energies compared to scattering on a heavy target, while leveraging existing detector technologies. Direct-detection experiments can also extend their reach to lower masses by exploiting the Migdal effect, where a nuclear recoil leads to electronic ionization or excitation. In this work, we combine these ideas to study the sensitivity of a hydrogen-doped LZ experiment (HydroX) and a future large-scale experiment such as XLZD. We find that HydroX could have sensitivity to dark matter masses below 10 MeV for both spin-independent and spin-dependent scattering, with XLZD extending that reach to lower cross sections. Notably, this technique substantially enhances the sensitivity of direct detection to spin-dependent proton scattering, well beyond the reach of any current experiments.

230

Dark Radiation Isocurvature from Cosmological Phase Transitions

Authors: Matthew Buckley¹; Peizhi Du¹; Nicolas Fernandez¹; Mitchell Weikert¹

¹ *Rutgers University*

Corresponding Author: peizhi.du@rutgers.edu

Cosmological first-order phase transitions are typically associated with physics beyond the Standard Model, and thus of great theoretical and observational interest. In this talk, I will show that a broad class of non-thermal first-order phase transitions could generate distinct large-scale isocurvature in dark radiation that can be observable in the CMB. We derive constraints on ΔN_{eff} from phase transitions based on CMB+BAO data, which can be much stronger than that from adiabatic initial conditions. I will also demonstrate that since perturbations of dark radiation have a non-Gaussian origin, searches for non-Gaussianity in the CMB could also place a stringent bound on ΔN_{eff} .

231

The Scotos side of Neutrinos

Authors: Kaladi Babu¹; Shreyashi Chakdar²; Vishnu Padmanabhan Kovilakam³

¹ *Oklahoma State University*

² *College of the Holy Cross*

³ *Universität Münster*

Corresponding Authors: kaladi.babu@okstate.edu, schakdar@holycross.edu, vpadmana@uni-muenster.de

We carry out a systematic investigation for minimal Scotogenic models based on a dark $U(1)_D$ gauge symmetry, in which the neutrino masses are induced at the one-loop level and include a chiral dark matter (DM) candidate. Assuming this $U(1)_D$ gauge symmetry is broken by only one Higgs singlet scalar that also generates masses to all dark fermions, we analyze the stability of the DM candidate which is ensured by a residual symmetry of $U(1)_D$ gauge symmetry. There can be different DM scenarios explored in this framework and we investigate the associated scalar and fermionic DM phenomenology of one of the minimal models.

232

Does the Sun Have a Dark Disk

Authors: Gustavo Alves¹; Mohammadreza Zakeri²; Pedro Machado³; Susan Gardner²

¹ *University Of Sao Paulo*

² *University of Kentucky*

³ *Fermilab*

Corresponding Authors: pmachado@fnal.gov, gardner@pa.uky.edu, m.zakeri@uky.edu, gustavo.figueiredo.alves@usp.br

The Sun's oblateness has been measured using optical observations. Its gravitational quadrupole moment has been deduced through helioseismology and measurements of its gravitational effects on Mercury's orbit. The distribution of mass within Mercury's orbit would only impact the orbital determination, suggesting that discrepancies among various types of assessments may indicate the possible existence of a non-luminous mass. For the first time, we have developed a method to combine these differing measurements to yield new, highly sensitive constraints on the mass distribution within Mercury's orbit. In this talk, we will show that the most precise measurements indicate the existence of a non-luminous disk within Mercury's orbit, with a mass significantly heavier than the modeled mass for the circumsolar dust ring observed by the Solar TErrestrial RElations Observatory (STEREO) mission. This suggests a substantial dark matter contribution. Furthermore, the long-standing inconsistency between the element abundances determined from the spectroscopy of the Sun's surface, and those inferred from its interior through helioseismology, can be reconciled if the Sun formed within a protoplanetary disk. We will discuss how our findings limit the presence of a dark disk or a spherical halo near the Sun and highlight the potential of future orbital measurements of Mercury and near-Sun asteroids to refine these constraints further.

233

Probing Heavy Neutral Lepton at Muon Collider and Beamdump Experiments

Author: Kunfeng Lyu¹

¹ *University of Minnesota*

Corresponding Author: lyu00145@umn.edu

Broad classes of solutions to the neutrino puzzles can be best tested by seeking the partners of SM light neutrinos, dubbed as heavy neutral leptons (HNLs). It can be parameterized by the HNL mass m_N and the mixing angle U_1 with the SM neutrino. In this talk, I will mainly discuss about probing two parameter space regime. One is for HNL mass larger than $O(100)$ GeV which can be tested at future high energy muon collider. Alternatively, if the HNL is lighter than the muon, it can be produced by the decays of muons and pions. The LSND experiment and the future beam dump experiment such as PIP2-BD can be exploited to improve the sensitivity.

234

Constraining non-standard neutrino interactions with neutral current events at long-baseline oscillation experiments

Author: Julia Gehrlein¹

¹ *Colorado State University*

Corresponding Author: julia.gehrlein@colostate.edu

Neutrino non-standard interactions (NSI) provide an attractive framework to study new physics in the neutrino sector in scattering experiments and in oscillation experiments where NSI lead to

new matter effects. At long-baseline oscillation experiments, so far, constraints on NSI have been derived using charged current events. In this manuscript we explore a new channel, namely neutral-current events at long-baseline experiments to constrain vector and axial-vector NSI. We introduce a framework to parametrize the effect of NSI on the cross section and then focus, as an example, on the NOvA experiment to derive constraints on NSI using real data.

235

Embedding the Universal See-Saw Mechanism in Pati-Salam Model

Authors: Kaladi S. Babu¹; Sumit Biswas¹

¹ *Oklahoma State University*

Corresponding Authors: kaladi.babu@okstate.edu, sumit.biswas@okstate.edu

Here we explore the integration of universal see-saw mechanism within the parity-symmetric Pati-Salam framework to address fermion mass generation. Incorporating vector-like fermions and a simple Higgs sector helps realize the hierarchical mass for all generations in this unified quark-lepton model. The neutrino sector seems pretty intriguing with the lightest neutrino acquiring mass through one-loop radiative corrections. With softly broken parity, we also see the possibility of addressing the strong CP problem in this setup.

236

Asymptotically free E_6 GUT and the generation of neutrino mass

Author: Vasja Susic¹

Co-authors: Borut Bajc ; K.S Babu ²

¹ *LNF, INFN*

² *Oklahoma State University*

Corresponding Authors: vasja.susic@lnf.infn.it, borut.bajc@ijs.si, kaladi.babu@okstate.edu

Grand Unified Theories (GUTs) represent an attractive possibility of physics beyond the Standard Model, with implications for both proton decay and neutrinos. The most studied examples are based on unified groups SU(5) and SO(10), while the exceptional case E_6 remains less understood.

In an asymptotically free E_6 GUT, model building is essentially limited to the use of irreducible representations of dimension 27 and 78 (the fundamental and the adjoint). I will present ongoing work in non-supersymmetric models regarding the determination of the necessary ingredients for both GUT symmetry breaking and a realistic Yukawa sector. Special attention will be given to the peculiarities associated with neutrino mass generation in this type of models.

237

Thermal Axion and Dark Radiation

Author: Fazlollah Hajkarim¹

¹ *University of Oklahoma*

Corresponding Author: fazlollah.hajkarim@ou.edu

We provide a comprehensive analysis of thermal axion production in the early universe, focusing on the KSVZ and DFSZ models. We extend our calculations to incorporate multiple mass thresholds, including the QCD phase transition, and provide a continuous production rate across these regimes. Employing updated cosmological data, we refine constraints on axion mass and examine the implications for dark radiation quantified as an effective number of additional neutrino species (ΔN_{eff}). Our rigorous approach revisits traditional approximations in axion production, highlighting the importance of precise calculations in anticipation of future CMB surveys and large scale structure observations. Also, we examine the validity of different computational approaches for the calculation of effective number of additional neutrino species.

238

E6 unification: intermediate symmetries, fermion masses and proton decay

Authors: Kaladi Babu^{None}; Borut Bajc^{None}; Vasja Susič^{None}

I will present a non-supersymmetric E6 GUT with the scalar sector consisting of $650 + 351' + 27$ dimensional representations. The intermediate symmetries which turn out to be realistic under the extended survival hypothesis (with minimal fine-tuning) are trinification symmetry $SU(3)_C \times SU(3)_L \times SU(3)_R$ with either LR or CR parity, and $SU(6)_{CR} \times SU(2)_L$. This means among others that they can reproduce correctly the light charged fermion and neutrino masses as well as the CKM and PMNS mixing matrices. Although the successful cases give a large range for proton lifetime estimates, all of them include regions consistent with current experimental bounds and within reach of forthcoming experiments.

239

Constraining Dwarf Galaxy Dark Matter Distributions: Spherical Jeans Analyses for Line-of-Sight and 3D Velocity Data

Author: Isabelle Goldstein¹

Co-author: Louie Strigari¹

¹ *Texas A&M University*

Corresponding Authors: isgoldstein@tamu.edu, strigari@tamu.edu

The stellar kinematics in dwarf galaxies can provide a wealth of information about its underlying dark matter distribution. Using line of sight velocity measurements for six classical dwarf galaxies of the Milky Way, we study whether ultralight bosonic dark matter is consistent with the gravitational potential extracted from stellar kinematics. It shows that axion-like particles with masses of order $m \sim 10^{-22}$ eV are inconsistent with the potential distribution in classical dwarf galaxies unless the hierarchical assembly of the Milky Way did not trace the mean evolution of Milky Way size halos. We also explore the use of three dimensional velocity measurements from Gaia data in spherical Jeans analyses to further constrain dark matter distributions.

240

-

Author: Brian Batell¹

¹ *University of Pittsburgh*

Corresponding Author: batell@pitt.edu

-

241

Probing Neutrinophilic Dark Matter: From Colliders to Supernovae

Author: Douglas Tuckler¹

¹ *TRIUMF and Simon Fraser University*

Corresponding Author: dtuckler@triumf.ca

New beyond-the-Standard Model mediators that couple predominantly to neutrinos are not yet probed by existing experimental searches. Such a neutrinophilic mediator is well motivated for addressing the origin of several neutrino-portal dark matter candidates, including thermal freeze-out and sterile-neutrino dark matter scenarios. In this talk, we explore the sensitivity to this scenario from two different approaches. In the first part of the talk we will explore the potential of the Forward Physics Facility (FPF) using the so-called “mono-neutrino signature”: neutrino charged-current scattering events associated with large missing transverse momentum, and excessive apparent tau-neutrino events. We will show that with this smoking-gun signature, the FPF has excellent sensitivity to probe this model in new regions of parameter space. In the second part of the talk we focus on astrophysical constraints on sterile neutrino DM, namely from core-collapse supernovae. Production and emission of DM can result in excessive energy loss of the supernova, leading to additional cooling. We will show that supernova cooling can constrain new regions of parameter space that complements terrestrial and cosmological probes. The results of this work present a nice complementarity among the collider and astrophysical frontiers.

242

Neutrino Interactions/TBD

Author: Vishvas Pandey¹

¹ *Fermilab*

Corresponding Author: vpandey@fnal.gov

Neutrino Interactions/TBD

243

Decoding the Flavor Evolution of Supernova Neutrinos: Shedding Light on Neutrino Electromagnetic Properties and Supernova Dynamics

Author: Yago Porto¹

¹ *ABC Federal University*

Corresponding Author: yago.porto@ufabc.edu.br

Two of the most important questions in neutrino astrophysics are: A) How can the extreme supernova environments, through which neutrinos propagate, be used to shed light on the new physics behind neutrino masses and mixing? B) What is the flavor evolution of supernova neutrinos in the latest stages of core collapse? In this talk, I will summarize our recent developments in these directions, emphasizing how the unique magnetic fields of supernova progenitors can modify neutrino evolution in the presence of both Dirac and Majorana neutrino magnetic moments during the first few tens of milliseconds of neutrino emission. For the rest of the neutrino emission (roughly 10 seconds), I will describe how standard matter effects can define the main features of the neutrino signal in terrestrial detectors despite the exotic neutrino-neutrino refraction that dominates in the supernova core.

244

TBD

Author: Tao Xu¹

¹ *The University of Oklahoma*

Corresponding Author: tao.xu@ou.edu

Talk details will be updated at a later date.

245

Quasi (Pseudo)-Dirac neutrinos

Author: Chee Sheng Fong¹

¹ *Universidade Federal do ABC*

Corresponding Author: sheng.fong@ufabc.edu.br

Here we will consider quasi (pseudo)-Dirac neutrinos

246

Dark Matter Raining on DUNE and Other Large Volume Detectors

Authors: Javier Acevedo¹; Joshua Berger²; Peter Denton³

¹ *SLAC National Accelerator Laboratory*

² *Colorado State University*

³ *Brookhaven National Laboratory*

Corresponding Authors: jfacev@slac.stanford.edu, joshua.berger@colostate.edu, peterbd1@gmail.com

Direct detection is a powerful means of searching for particle physics evidence of dark matter (DM) heavier than about a GeV with $\mathcal{O}(\text{ton})$ volume, low-threshold detectors.

In many scenarios, some fraction of the DM may be boosted to large velocities enhancing and generally modifying possible detection signatures. We investigate the scenario where 100% of the DM may be boosted at the Earth due to new attractive long-range forces. This opens up two main improvements in detection capabilities: 1) the detection signatures are stronger opening up large-volume neutrino detectors, such as DUNE, Super-K, Hyper-K, and JUNO, as possible DM detectors, and 2) the large boost allows for detectable signatures of sub-GeV DM. At lower boosts, a modified, higher-than-usual energy signal could be accessible at direct detection experiments such as LZ. In addition, the model leads to a significant anisotropy in the signal with the DM flowing dominantly vertically at the Earth's surface instead of the typical approximately isotropic DM signal. We develop the theory behind this model and also calculate realistic constraints using a detailed GENIE simulation of the signal inside detectors.

247

Cosmological Stasis and Its Realization from Dynamical Scalars

Author: Brooks Thomas¹

Co-authors: Fei Huang²; Keith R. Dienes³; Lucien Heurtier⁴; Timothy M. P. Tait⁵

¹ *Lafayette College*

² *Weizmann Institute*

³ *University of Arizona*

⁴ *King's College London*

⁵ *University of California Irvine*

Corresponding Authors: ttait@uci.edu, lucien.heurtier@kcl.ac.uk, fei.huang@weizmann.ac.il, thomasbd@lafayette.edu, dienes@arizona.edu

It has recently been realized that many extensions of the Standard Model give rise to cosmological histories exhibiting extended epochs of cosmological stasis —epochs wherein the abundances of multiple energy components (such as matter, radiation, or vacuum energy) remain effectively constant despite cosmological expansion. The emergence of a stasis epoch is not a consequence of fine-tuning in cosmologies of this sort; rather, stasis turns out to be a global attractor toward which the universe naturally evolves for a broad range of initial conditions. In this talk, I shall review the general conditions under which stasis emerges in such scenarios and explore some of its potential implications for cosmological observables such as the matter power spectrum and the stochastic gravitational-wave background. I shall also discuss a particular realization of stasis involving a collection of scalar fields, each of which dynamically transitions from a period of slow roll to a period of rapid oscillation around its potential minimum as the universe expands. As I shall demonstrate, not only does cosmological stasis arise in such scenarios, but the system of dynamical scalars also exhibits novel features not seen in previous realizations of stasis, including a tracking behavior wherein the effective equation of state for the universe as a whole evolves toward the equation of state of this energy component. The emergence of such tracking behavior has potential model-building implications in the context of dark-energy and cosmic-inflation scenarios.

248

Neutrino and light dark matter physics with directional detectors

Author: Diego Aristizabal¹

¹ *USM*

Corresponding Author: diego.aristizabal@usm.cl

In this talk I will review a few aspects of Coherent Elastic neutrino-Nucleus Scattering (CEvNS) and light dark matter (LDM) production in meson decays. In particular, I will focus on potential measurements using neutrino beams at FNAL. I will show that, combined with directional detectors, these beams offer an avenue for CEvNS measurements as well as for LDM detection. Such measurements will provide complementary information to that coming from other CEvNS facilities as well as from other non-directional detectors looking for LDM.

249

Direct Collapse Black Holes from Dark Matter Annihilation

Author: Flip Tanedo¹

¹ *UC Riverside*

Corresponding Author: flip.tanedo@ucr.edu

Pre-stellar galactic halos are highly sensitive to soft radiation: the presence of sources of $O(10 \text{ eV})$ Lyman-Werner radiation changes the gas chemistry and prevent the standard fragmentation of the gas. Rather than producing Population III stars, this may instead lead to direct collapse black holes. Observations of supermassive black holes at high redshift have long been suspected to be evidence for direct collapse black holes. Recent studies have explored the possibility that direct collapse may be influenced by new particle physics.

We present a simple dark matter model where resonant annihilation can dissociate molecular hydrogen and discuss the assumptions that are necessary to induce direct collapse black holes. In these models, $O(10 \text{ MeV})$ dark matter annihilates into electron-positron pairs which produce Lyman-Werner radiation by inverse Compton scattering CMB light. We present a self-consistent modeling of H_2 self-shielding that highlights the challenges when building models for direct collapse.

250

Probing self-interacting sterile neutrino dark matter with the diffuse supernova neutrino background

Author: Anna Suliga¹

Co-authors: Anupam Ray²; Baha Balantekin³; George Fuller⁴

¹ *UC Berkeley*

² *UC, Berkeley*

³ *UW Madison*

⁴ *UC, San Diego*

Corresponding Authors: anupam.ray@berkeley.edu, gfuller@physics.ucsd.edu, baha@physics.wisc.edu, asuliga@berkeley.edu

The neutrinos in the diffuse supernova neutrino background (DSNB) travel over cosmological distances and this provides them with an excellent opportunity to interact with dark relics. We show that a cosmologically-significant relic population of keV-mass sterile neutrinos with strong self-interactions could imprint their presence in the DSNB. The signatures of the self-interactions would be “dips” in the otherwise smooth DSNB spectrum. Upcoming large-scale neutrino detectors, for example Hyper-Kamiokande, have a good chance of detecting the DSNB and these dips. If no dips are detected, this method serves as an independent constraint on the sterile neutrino self-interaction strength and mixing with active neutrinos. We show that relic sterile neutrino parameters that evade X-ray and structure bounds may nevertheless be testable by future detectors like TRISTAN, but may

also produce dips in the DSNB which could be detectable. Such a detection would suggest the existence of a cosmologically-significant, strongly self-interacting sterile neutrino background, likely embedded in a richer dark sector.

251

Planting Native Plants at Sanford Lab Homestake Visitor Center

252

Communicating with Media (Speakers: Ann Melti and Mike Ray, SDSTA/SURF)

Corresponding Author: ametli@sanfordlab.org

253

Communicating with Media - 3 Minute Challenge (Ann Melti and Mike Ray - SDSTA/SURF)

254

Career Panel

255

Scientist Mentor/Student Speed Match Up and Shadow

256

Communicating with Media - Ann Melti and Mike Ray (SDSTA/SURF)

Corresponding Author: ametli@sanfordlab.org

257

3 Minute Challenge - Ann Melti and Mike Ray (SDSTA/SURF)

258

Jet-SIFTing for Dark Sector Physics in a Hidden Valley

Author: Joel Walker¹

¹ *Sam Houston State University*

Corresponding Author: jww004@shsu.edu

We consider a Hidden Valley model which generates showering from strong dynamics within the dark sector followed by decays back into Standard Model states. Our interest is the limit of smaller dark pion masses, which create a high multiplicity of final states. The reconstruction of dark sector masses in such a setting is obscured by a thick combinatoric background. We apply the new SIFT (Scale-Invariant Filtered Tree) jet clustering algorithm to the reconstruction of simulated events of this type. By cutting an ordered slice through possible recombinations, the SIFT algorithm may help lift backgrounds of the described variety.

259

Skimming tau neutrinos and optical Cherenkov signals

Author: Mary Hall Reno¹

¹ *University of Iowa*

Corresponding Author: mary-hall-reno@uiowa.edu

Optical Cherenkov signals from up-going air showers show promise for detection of skimming tau neutrinos. We review the theoretical inputs to evaluating the sensitivities to transient neutrino sources of orbital and sub-orbital Cherenkov telescopes.

260

Detection of cosmogenic (B)SM signals with nuclear inelastic scattering

Author: Doojin Kim¹

¹ *Texas A&M University*

Corresponding Author: doojin.kim@tamu.edu

I will discuss the detection of sub-GeV-range cosmogenic signals at large-volume neutrino detectors such as DUNE, SK/HY, and JUNO, utilizing nuclear inelastic scattering channels featuring nuclear deexcitation gamma-ray lines. I will first briefly discuss the detection of neutrino signals and point out the potential of observing the 13-MeV oxygen line at SK. I will then propose a new approach to search for light dark matter (DM) in the range of keV-GeV in the context of cosmic-ray boosted dark matter. I will show that using a hadrophilic dark-gauge-boson-portal model as a benchmark, the nuclear inelastic channels generally provide better sensitivity than the elastic scattering for a large region of light DM parameter space.

261

BSM Physics with Gravitational Wave Detectors

Author: Kuver Sinha¹

¹ *University of Oklahoma*

Corresponding Author: kuver.sinha@ou.edu

Future gravitational wave detectors probing the mHz - nHz frequency range will provide a unique opportunity for BSM physicists to study new physics. Neutron star and white dwarf mergers can serve as axion probes, while extreme mass ratio inspirals can constrain dark forces. Gravitational wave detectors will also probe early first order phase transitions. I will discuss some ongoing work and future ideas in these directions.

262

eV Sterile Neutrino Paradox

Authors: Alexei Yu. Smirnov^{None}; Lucas Puetter¹; Sudip Jana^{None}

¹ *Max-Planck-Institut für Kernphysik*

Corresponding Authors: sudip.jana@okstate.edu, smirnov@mpi-hd.mpg.de, lucas.puetter@mpi-hd.mpg.de

Light sterile neutrinos are frequently featured in various theories, particularly to explain short-baseline anomalies or as candidates for dark matter. These scenarios necessitate non-zero mixing with active neutrinos, often disregarding the contributions to the active neutrino mass matrix. We demonstrate that this mixing induces contributions to the active neutrino mass matrix, thereby affecting neutrino oscillation results. We conduct a rigorous analysis and present stringent bounds on these induced masses for each entry of a 3×3 active neutrino mass matrix. Furthermore, we examine whether the induced matrix can elucidate the distinctive pattern of lepton mixing and the neutrino mass spectrum, while discussing the various phenomenological implications.

263

CETUP* Workshop Overview

264

Discussion and Collaboration

265

TALK: New Constraints on Neutrino-Dark Matter Interactions

Corresponding Author: bdev@wustl.edu

Author: Bhupal Dev

We present a comprehensive analysis of nonstandard neutrino interactions with the dark sector in an effective field theory framework. We implement a full catalog of constraints on the parameter

space of the neutrino-dark matter/mediator couplings and masses, including bounds coming from cosmology, astrophysics, as well as new laboratory constraints, such as from invisible Z decays and rare meson decays. We find that most of the benchmarks in the dark matter mass-coupling plane adopted in previous studies to get an observable effect are actually ruled out by a combination of these constraints. Finally, as an application of our results, we consider the case of galactic supernova neutrinos, identify new benchmark points for future observational prospects of the attenuation of the galactic supernova neutrino flux, compute the full set of cascade equations and sky maps for different dark matter density profiles in the Galaxy, and comment on their implications for the detection prospects in future large-volume neutrino experiments such as DUNE, Hyper-K and JUNO.

266

TALK: TBD (open slot available)

267

Shuttle, Travel to Ross

268

Does the Sun have a dark disk?

Authors: Gustavo Alves¹; Susan Gardner²; Pedro Machado³; Mohammadreza Zakeri²

¹ *University Of Sao Paulo*

² *University of Kentucky*

³ *Fermilab*

Corresponding Authors: gustavo.figueiredo.alves@usp.br, m.zakeri@uky.edu, pmachado@fnal.gov, gardner@pa.uky.edu

The nature of dark matter is unknown, as are its total mass and distribution within the Solar system, though constraints from measurements of planetary orbits exist. I discuss how differing measurements of the Sun's gravitational quadrupole moment can combine to yield new and highly sensitive constraints on the mass distribution within Mercury's orbit. The best determinations point with high confidence to the existence of a non-luminous disk coplanar with Mercury's orbit, and the mass estimates associated with known matter, although uncertain, point to a significant dark-matter contribution, for which macroscopic dark-matter models may suit best.

269

TALK: Direct Collapse Black Holes from Dark Matter Annihilation

Corresponding Author: flip.tanedo@ucr.edu

Pre-stellar galactic halos are highly sensitive to soft radiation: the presence of sources of $O(10 \text{ eV})$ Lyman-Werner radiation changes the gas chemistry and prevent the standard fragmentation of the gas. Rather than producing Population III stars, this may instead lead to direct collapse black holes. Observations of supermassive black holes at high redshift have long been suspected to be evidence

for direct collapse black holes. Recent studies have explored the possibility that direct collapse may be influenced by new particle physics.

We present a simple dark matter model where resonant annihilation can dissociate molecular hydrogen and discuss the assumptions that are necessary to induce direct collapse black holes. In these models, $O(10 \text{ MeV})$ dark matter annihilates into electron-positron pairs which produce Lyman-Werner radiation by inverse Compton scattering CMB light. We present a self-consistent modeling of H_2 self-shielding that highlights the challenges when building models for direct collapse.

270

TALK: TBD (open slot available)

271

Discussions, Collaboration and Lunch

272

Current Challenges and Opportunities in Physics Community (open discussion)

Corresponding Author: barbara.szczerbinska@tamucc.edu

Academia
Research
Funding
Promotion and tenure
Mentoring

273

Global Scientist Social

Corresponding Author: rmcgehee@umn.edu

274

SURF Underground Tour

275

Discussion and Collaboration; Mentor/Student Shadow

276

A SURF Low Background Module (SLoMo)

Author: Juergen Reichenbacher¹

¹ *South Dakota School of Mines and Technology*

Corresponding Author: juergen.reichenbacher@sdsmt.edu

With radiopurity controls and targeted design modifications a kton-scale liquid argon time projection chamber similar to DUNE could be used for enhanced low energy physics searches. This includes improved sensitivity to supernova and solar neutrinos, and other rare event searches, such as WIMP dark matter, while simultaneously serving long-baseline neutrino physics. This talk will present simulation studies to evaluate physics sensitivities of such a module. It will also discuss R&D to develop large-scale radiopurity controls necessary to construct such a detector.

277

The Institute for Underground Science at SURF Overview and Update (room 132)

278

The Institute for Underground Science at SURF Overview and Update (room 132)

279

The Institute for Underground Science at SURF Overview and Update (room 132)

280

REMINDER NOTE ONLY: please make sure to bring a brown bag lunch for today

281

TALK: Detection of cosmogenic (B)SM signals with nuclear inelastic scattering

Corresponding Author: doojin.kim@tamu.edu

Author: Doojin Kim

I will discuss the detection of sub-GeV-range cosmogenic signals at large-volume neutrino detectors

such as DUNE, SK/HY, and JUNO, utilizing nuclear inelastic scattering channels featuring nuclear deexcitation gamma-ray lines. I will first briefly discuss the detection of neutrino signals and point out the potential of observing the 13-MeV oxygen line at SK. I will then propose a new approach to search for light dark matter (DM) in the range of keV-GeV in the context of cosmic-ray boosted dark matter. I will show that using a hadrophilic dark-gauge-boson-portal model as a benchmark, the nuclear inelastic channels generally provide better sensitivity than the elastic scattering for a large region of light DM parameter space.